


令和6年度 橘湾・椿泊湾 有害プランクトン調査（5月21日調査）徳島県水産研究課

St.	採水時刻	採水層 (m)	水温 (°C)	塩分	透明度 (m)	細胞/mL			
						シャットネラ 区分不能	シャットネラ オバータ	カレニア ミキモトイ	コクロディニウム ポリクリコイデス
1	09:48	0-5			6.1	-	-	-	-
		0	18.59	31.66		-	-	-	-
		1	18.52	31.67					
		5	17.88	32.08					
		10	17.37	32.53					
		B-1	17.37	32.51					
2	09:38	0-5			4.3	-	-	-	-
		0	19.07	30.84					
		1	18.69	31.46					
		5	17.58	32.30					
		B-1	17.40	32.42					
3	09:26	0-5			4.5	-	-	-	-
		0	18.89	31.85		-	-	-	-
		1	18.85	31.85					
		5	18.37	32.11					
		B-1	18.32	32.13					
3'	09:20	0-5			5.1	-	-	-	-
		0	19.01	31.82					
		1	18.81	31.82					
		5	18.18	32.17					
		B-1	17.91	32.31					
4	09:05	0-5			6.0	-	-	-	-
		0	18.46	32.04					
		1	18.42	31.98					
		5	18.29	32.02					
		10	18.15	32.17					
		15	17.26	32.65					
		B-1	17.19	32.89					



備 考

・珪藻は橘湾はスケルトネマ属主体、椿泊湾はキートセロス属主体で、前回調査と同程度でした。

珪藻細胞数 橘湾St.1(0-5m):634 cells/ml
椿泊湾St.3(0-5m):461 cells/ml

令和6年度 橘湾・椿泊湾 有害プランクトン調査（5月21日調査）徳島県水産研究課

St.	採水時刻	採水層 (m)	水温 (℃)	塩分	透明度 (m)	細胞/mL				備 考
						シャットネラ 区分不能	シャットネラ オーバータ	カレニア ミキモトイ	コクロディニウム ポリクリコイデス	
1	08:27	0-5			5.9	-	-	-	-	
		0	18.37	31.55						
		1	18.02	31.94						
		5	17.43	32.40						
		B-1	17.25	32.67						
2	08:35	0-5			5.8	-	-	-	-	
		0	18.13	31.66		-	-	-	-	
		1	17.82	32.04						
		5	17.39	32.46						
		10	17.29	32.78						
		15	17.26	33.03						
		B-1	17.25	33.03						
3	10:13	0-5			5.5	-	-	-	-	
		0	18.47	31.74						
		1	18.17	31.87						
		5	17.78	32.14						
		10	17.52	32.39						
		15	17.32	32.72						
		B-1	17.25	33.03						
4	08:44	0-5			8.0	-	-	-	-	
		0	18.22	31.58						
		1	17.86	31.74						
		5	17.69	32.26						
		10	17.49	32.43						
		15	17.35	32.65						
		20	17.30	33.09						
		B-1	17.32	33.13						