


令和6年度 ウチノ海環境調査（12月19日調査）徳島県水産研究課

St.	採水時刻	採水層 (m)	水温 (°C)	塩分	透明度 (m)	DO (mg/L)	細胞/L			
							ユーカンピア ゾディアクス	コシノディスカス ワイレシー	コシノディスカス spp.	
1	08:56	0-5			8.0		-	50	10	
		0	14.51	32.28		7.9				
		1	14.51	32.30		7.9				
		5	14.53	32.32		7.9				
		10								
		B-1	14.53	32.35		7.9				
2	09:01	0-5			7.1		-	10	20	
		0	14.02	32.32		8.1				
		1	14.04	32.34		8.1				
		5	14.01	32.33		8.1				
		B-1	13.95	32.33		8.1				
3'	09:07	0-5			7.0		-	30	-	
		0	14.23	32.29		8.1	-	20	-	
		1	14.25	32.34		8.1	-	-	20	
		5	14.15	32.31		8.1	0.14	-	50	
		10	14.08	32.33		8.1				
		B-1	14.08	32.33		8.1	-	20	20	
4	09:14	0-5			7.6		-	-	-	備 考 ・藻類を色落ちさせる珪藻であるコシノディスカス属及びユーカンピア属が検出されましたが低密度でした。 ・珪藻は前回調査時より増加しましたが低密度でした。 珪藻細胞数: 7 cells/ml (St.5 0-5m)
		0	14.47	32.27		8.0				
		1	14.53	32.34		8.0				
		5	14.38	32.32		8.0				
		10	14.23	32.31		8.0				
		B-1	14.18	32.31		8.0				
5	09:20	0-5			5.9		-	-	10	
		0	13.93	32.28		8.2				
		1	13.94	32.29		8.2	-	30	30	
		5	13.94	32.29		8.1	0.01	50	-	
		10	13.94	32.29		8.1	-	-	10	
		B-1	13.93	32.29		8.1				
6	09:29	0-5			7.0		-	10	30	
		0	13.72	32.27		8.2				
		1	13.72	32.28		8.2				
		5	13.74	32.27		8.2				
		10	13.72	32.27		8.1				
		B-1	13.71	32.27		8.1				