


令和6年度 ウチノ海環境調査（3月19日調査）徳島県水産研究課

St.	採水時刻	採水層 (m)	水温 (℃)	塩分	透明度 (m)	DO (mg/L)	細胞/ml	細胞/l	
							ユーカンピア ゾディアクス	コシノディスカス ワイレシー	コシノディスカス spp.
1	08:58	0-5			9.0		-	-	-
		0	9.14	32.49		9.3			
		1	9.14	32.49		9.3			
		5	9.14	32.51		9.3			
		10	9.14	32.54		9.3			
		B-1	9.14	32.53		9.3			
2	09:03	0-5			8.0		-	-	-
		0	9.24	32.44		9.5			
		1	9.30	32.44		9.5			
		5	9.26	32.44		9.5			
		B-1	9.14	32.44		9.4			
3'	09:08	0-5			8.0		-	-	10
		0	9.28	32.43		9.3	-	-	-
		1	9.31	32.46		9.3	-	-	-
		5	9.29	32.47		9.3	-	-	-
		10	9.30	32.51		9.2			
		B-1	9.32	32.50		9.2	-	-	-
4	09:14	0-5			8.0		-	-	-
		0	9.27	32.44		9.4			
		1	9.28	32.45		9.4			
		5	9.28	32.47		9.3			
		10	9.29	32.49		9.3			
		B-1	9.29	32.50		9.2			
5	09:19	0-5			8.5		-	-	-
		0	9.16	32.41		9.4			
		1	9.16	32.42		9.4			
		5	9.16	32.42		9.4			
		10	9.15	32.42		9.4	0.04	-	-
		B-1	9.15	32.42		9.4			
6	09:28	0-5			8.5		-	-	-
		0	9.20	32.38		9.5			
		1	9.21	32.38		9.5			
		5	9.21	32.37		9.4			
		10	9.21	32.38		9.4			
		B-1	9.21	32.32		9.4			



備考

・藻類を色落ちさせる珪藻のコシノディスカス属及びユーカンピア属が検出されましたが、低密度でした。

・珪藻は非常に低密度で前回調査時から減少しました。

珪藻細胞密度 St.5(0-5m): 9 cells/ml