


令和6年度 北灘 環境調査（3月10日調査）徳島県水産研究課

St.	採水時刻	採水層 (m)	水温 (℃)	塩分	透明度 (m)	DO (mg/L)	細胞/L		細胞/mL	
							コシノディスカス ワイレシー	コシノディスカス spp.	ユーカンピア ゾディアクス	
2	09:40	0-5			10.8		-	-	-	
		0	8.47	32.30		9.8				
		1	8.37	32.30		9.8	-	-	-	
		5	8.27	32.29		9.8	10	-	-	
		10	8.15	32.30		9.8	-	-	-	
		15	8.16	32.31		9.7				
		20	8.13	32.32		9.5				
		B-1	8.18	32.34		9.4	-	-	-	
4	09:08	0-5			10.1		-	-	-	
		0	8.84	32.33		9.8	-	-	-	
		1	8.82	32.40		9.8	-	-	-	
		5	8.82	32.43		9.7	-	-	-	
		10	8.56	32.40		9.7	-	-	-	
		15	8.50	32.45		9.6				
		20	8.56	32.50		9.5	-	-	-	
		25	8.65	32.50		9.4				
		30	8.65	32.51		9.4	-	-	-	
		35	8.64	32.51		9.4				
		40	8.64	32.52		9.3				
		B-1	8.64	32.52		9.3	-	-	-	
5	10:09	0-5			12.8		-	-	-	備 考 ・藻類を色落ちさせる珪藻について、コシノディスカス属が検出されましたが低密度でした。 ・珪藻は非常に低密度でした。 珪藻細胞数 St.4(0-5m): 4 cells/ml
		0	8.68	32.35		9.9				
		1	8.61	32.29		9.9	-	-	-	
		5	8.32	32.33		9.9	-	-	-	
		10	8.26	32.35		9.8	-	-	-	
		15	8.39	32.40		9.8				
		20	8.65	32.49		9.5				
		25	8.65	32.50		9.4				
		30	8.66	32.52		9.3				
		B-1	8.67	32.52		9.3	10	-	-	