


令和7年度 ウチノ海環境調査（6月4日調査）徳島県水産研究課

St.	採水時刻	採水層 (m)	水温 (°C)	塩分	透明度 (m)	カレニア ミキモトイ	
1	09:08	0-5			4.5	-	備 考 ・有害種は検出されませんでした。 ・珪藻はシュードニッチア属主体で低密度でした。 珪藻細胞密度 : 1269cells/ml (St.5 0-5m)
		0	18.28	32.32			
		1	18.29	32.34			
		5	18.20	32.37			
		10	18.14	32.40			
		B-1	18.16	32.40			
2	09:12	0-5			5.1	-	
		0	18.45	32.26			
		1	18.33	32.32			
		5	18.09	32.45			
		B-1	18.08	32.46			
3'	09:17	0-5			5.1	-	
		0	18.54	32.31			
		1	18.45	32.35			
		5	18.13	32.41			
		10	18.08	32.42			
		B-1	18.08	32.42			
4	09:23	0-5			5.9	-	
		0	18.79	32.24			
		1	18.57	32.30			
		5	18.21	32.45			
		10	18.10	32.48			
		B-1	18.08	32.49			
5	09:28	0-5			5.0	-	
		0	18.79	32.16			
		1	18.78	32.16			
		5	18.35	32.38			
		10	17.95	32.48			
		B-1	17.96	32.48			
6	09:36	0-5			5.0	-	
		0	18.88	32.05			
		1	18.86	32.06			
		5	18.65	32.16			
		10	17.99	32.46			
		B-1	17.95	32.47			